

Imaging the Surface States of a Strongly Correlated Topological Insulator, SmB₆

The poster features a blue background with white and yellow text. At the top left, it reads 'Colloquium Frontiers of Condensed Matter Physics' with a dedication to Prof. Nicolás Cabrera (1913-1989). The year '2018' is prominently displayed in a large white font. The date '12/04' is shown in a white box. The title of the talk is 'Imaging the Surface States of a Strongly Correlated Topological Insulator, SmB₆' by Jenny Hoffman. A cartoon illustration of Jenny Hoffman stands next to a chalkboard that contains the title and a diagram of a crystal lattice. A blue box on the right provides details: 'When: 12 April at 12h 30', 'Where: Sala de grados, módulo 08, Facultad de Ciencias, UAM', and a short abstract of the talk. At the bottom right, the 'Instituto Nicolás Cabrera' logo is visible.

INC COLLOQUIUM - OFFICIAL ANNOUNCEMENT

Title: Imaging the Surface States of a Strongly Correlated Topological Insulator, SmB₆.

When: 12 April, 2018, 12h30

Where: Sala de Grados, Módulo 08, Faculty of Science, UAM.

Speaker: Jenny Hoffman, Department of Physics, Faculty of Arts And Sciences, Harvard University, USA.

The prediction and subsequent discovery of robust spin-polarized surface states on topological band insulators has launched a new subfield of physics over the last decade. When topology is combined with strong electron correlations, even more interesting states of matter can arise, such as quantum anomalous Hall states, fractionalization, non-Abelian exchange statistics, and many-body localization that preserves the topology of the insulating state against thermal destruction. Here I will give a general introduction to topological materials, and show the first direct proof of a strongly correlated topological insulator. Using scanning tunneling microscopy to probe the real and momentum space structure of SmB₆, we quantify the opening of a Kondo insulating gap. Within that gap, we discover linearly dispersing surface states with effective masses reaching $m = (330 \pm 20)m_e$, the heaviest observed Dirac states in any material. The prodigious density of the Dirac states near zero energy magnifies both their susceptibility to novel orders, and their potential utility. Our observations present the first opportunity to explore a strongly correlated topological state of matter.

Quantum Transport in Topological Materials

XXIV International Summer School 'Nicolás Cabrera'

September 4-8, 2017

Miraflores de la Sierra

When: September 4-8, 2017

Where: Residencia La Cristalera, Miraflores de la Sierra, Madrid, Spain

Contacts:

Inés Ortiz - e-mail: school@nicolascabrera.es

Manuela Moreno - e-mail: manuela.moreno@uam.es

Organizers:

Eduardo Lee, IFIMAC, UAM.

Elsa Prada, IFIMAC, UAM.

Alfredo Levy Yeyati, IFIMAC, UAM

School Scopes and Goals

Topological materials constitute an exciting and very active research area in condensed matter physics. It studies new states of matter whose bulk properties are similar to those of 'ordinary' materials but that, at the same time, display edge or boundary states with very exotic properties. Since the discovery of topological insulators, roughly a decade ago, the field has rapidly expanded with the identification of other topological materials, such as topological superconductors and Weyl semimetals. This Summer School will gather leading international experts to provide an introduction to the basic concepts underlying topology in condensed matter systems, followed by a discussion of recent developments, with a focus on quantum transport and hybrid devices. The goal is to cover not only theoretical aspects, but to also address the experimental progress, including the detection and manipulation of states

associated with these materials.

School Topics

Hybrid devices (quantum dots, nanowires, heterostructures).

Topological insulators and superconductors.

Weyl semimetals.

Topological quantum computing.

Invited Speakers

Ramón Aguado (ICMM-CSIC, Madrid)

Alberto Cortijo (ICMM-CSIC, Madrid)

Silvano De Franceschi (CEA, Grenoble)

Reinhold Egger (Heinrich Heine Univ., Düsseldorf)

Klaus Ensslin (ETH, Zürich)

Claudia Felser (Max Planck Inst. for Chemical Physics of Solids, Dresden)

Marcelo Goffman (CEA, Saclay)

Sophie Guéron (Univ. Paris Sud, Orsay)

Jelena Klinovaja (Univ. Basel)

Leo Kouwenhoven (QuTech, Delft Univ. of Technology)

Rosa López (Univ. Baleares)

Fabrizio Nichele (Niels Bohr Institute, Copenhagen)

Yuval Oreg (Weizmann Inst. of Science)

Pablo San-Jose (ICMM-CSIC, Madrid)

Jörg Schäfer (Univ. Würzburg)

Patrik Recher (TU Braunschweig)

Shinsei Ryu (Univ. Chicago)

Felix von Oppen (Freie Univ. Berlin)

Organized with the collaboration of

The logo for Fundación BBVA, featuring the word "Fundación" in a blue sans-serif font and "BBVA" in a larger, bold, blue sans-serif font.

Further information:

Nicolás Cabrera Institute - [INC website](#)

XXIV International Summer School Nicolás Cabrera in the newspapers - [Atlasagencia](#)


Weyl Semimetals in 3D Optical Lattices, and Synthetic Gauge Fields in Strongly Interacting 1D Bose Gases

Title: Weyl Semimetals in 3D Optical Lattices, and Synthetic Gauge Fields in Strongly Interacting 1D Bose Gases.

When: Friday, November 25, (2016), 12:00.

Place: Departamento de Física Teórica de la Materia Condensada, Facultad Ciencias, Module 5, Seminar Room (5th Floor).

Speaker: Hrvoje Buljan, Department of Physics, Faculty of Science, University of Zagreb, Bijenička c. 32, 10000 Zagreb, Croatia.



We discuss two important directions of research of synthetic topological quantum matter [1]: (i) topological phases in 3D optical lattices, more specifically Weyl semimetals in ultracold atomic gases [2], and (ii) strongly interacting Bose gases in synthetic gauge fields [3]. Interacting Bose gases in synthetic magnetic fields [1] hold great potential for discovering and exploring novel topological states of matter [4], in a similar fashion as the fractional quantum Hall effect for electrons in strong magnetic fields. Here we investigate laser assisted tunneling in a strongly interacting one-dimensional Bose gas [3] (the Tonks-Girardeau gas [5]) in optical lattices. We find that the stroboscopic dynamics of the Tonks-Girardeau gas in a continuous Wannier-Star-ladder potential, supplemented with laser assisted tunneling, effectively realizes the ground state of one-dimensional hardcore bosons in a discrete lattice [6] with nontrivial hopping phases. Next, we show that, in three-dimensional optical lattices, laser assisted tunneling can be used for realizing a Hamiltonian with Weyl points [3]. Weyl points are synthetic magnetic monopoles that exhibit a robust, three-dimensional linear dispersion, identical to the energy-momentum relation for relativistic Weyl fermions [3], which are not yet discovered in particle physics. Relation to analogous studies in optics will be mentioned [8].

References

N. Goldman, G. Juzeliunas, P. Ohberg, I. B. Spielman, Rep. Prog. Phys. 77, 126401 (2014).

Tena Dubček, Colin J. Kennedy, Ling Lu, Wolfgang Ketterle, Marin Soljačić, Hrvoje Buljan, Weyl points in three-dimensional optical lattices: Synthetic magnetic monopoles in momentum space, Phys. Rev. Lett. 114, 225301 (2015).

Karlo Lelas, Nikola Drpić, Tena Dubček, Dario Jukić, Robert Pezer, and Hrvoje Buljan, Laser assisted tunneling in a Tonks-Girardeau gas, New J. of Phys. 18, 095002 (2016).

X. Chen, Z.-C. Gu, Z.-X. Liu, X.-G. Wen, arXiv:1301.0861 [cond-mat.str-el]; Science 338, 1604 (2012).

M. Rigol and A. Muramatsu, Phys. Rev. Lett. 93, 230404 (2004).

R. Pezer and H. Buljan, Phys. Rev. Lett. 98, 240403 (2007).

Jorge Bravo-Abad, Ling Lu, Liang Fu, Hrvoje Buljan and Marin Soljačić, Weyl points in photonic-crystal superlattices, 2D Mater. 2, 034013 (2015).

Tena Dubček, Karlo Lelas, Dario Jukić, Robert Pezer, Marin Soljačić and Hrvoje Buljan,
The Harper-Hofstadter Hamiltonian and conical diffraction in photonic lattices with
grating assisted tunneling, *New J. Phys.* 17, 125002 (2015).

[More information on IFIMAC Website](#)
